

Quark Masses from Gaugino Condensation in String Theories

Christos Kokorelis ^{1,*}, Van E. Mayes ^{2,†} and Dimitri V. Nanopoulos ^{2,3,4‡}

¹ *Institute of Nuclear Physics, N.C.S.R. Demokritos, 15310, Athens, Greece*

² *George P. and Cynthia W. Mitchell Institute for Fundamental Physics,
Texas A&M University, College Station, TX 77843, USA*

³ *Astroparticle Physics Group, Houston Advanced Research Center (HARC), Mitchell Campus, Woodlands, TX 77381, USA*

⁴ *Academy of Athens, Division of Natural Sciences,
28 Panepistimiou Avenue, Athens 10679, Greece*

We present a mechanism able to generate the perturbatively absent up/down $\langle \mathbf{10} \cdot \mathbf{10} \cdot \mathbf{5}^H \rangle$ quark Yukawa couplings of SU(5)/flipped SU(5) GUTS in Type II orientifold compactifications with D-branes. The mechanism works when there are Sp(N) gauge groups involved. The $\bar{\mathbf{5}}$'s get charged under the Sp(N) gauge groups and the generation of quark masses proceeds via the generation of the fermionic Sp(N) singlet condensate $\langle \bar{\mathbf{5}} \cdot \bar{\mathbf{5}} \cdot \bar{\mathbf{5}} \cdot \bar{\mathbf{5}} \rangle$ in the term $(1/M_s^5) \mathbf{10} \cdot \mathbf{10} \cdot \langle \bar{\mathbf{5}} \cdot \bar{\mathbf{5}} \cdot \bar{\mathbf{5}} \cdot \bar{\mathbf{5}} \rangle$. Also non-chiral states charged under Sp gauge groups may become constrained by the requirement of Sp's becoming strongly coupled.

I. INTRODUCTION

Non-perturbative mechanisms are very important in modern particle theories as they are used to solve various problems. For example, QCD instantons [1] are a famous solution in QCD for solving the η' -meson mass problem [2] and SU(2) instantons in generating baryon number violating interactions. On the other hand in string compactifications worldsheet instantons have been associated with genus zero instantons that generate corrections to the spacetime superpotential [3]. Moreover gaugino condensation has been connected to the formation of hidden matter condensates [4] and supersymmetry breaking. Also in the context of intersecting brane worlds [5, 6] hidden sector confinement has been utilized to decouple hidden sector states at high energies [7]. On the other hand as intersecting brane models (IBM) [5, 6] fix some of the moduli either via supersymmetry conditions in intersections or fluxes, it is quite desirable to find a flipped SU(5) (FSU(5)) IBM string model which could mimic the phenomenological success of fermionic FSU(5) [8].

Quite recently, in the context of type II compactifications, D2-brane (E2) instantons wrapping special Lagrangian three-cycles of the internal Calabi-Yau [9] can generate perturbatively absent couplings like Majorana terms for right handed neutrinos, μ -terms in the MSSM [9, 10, 11] or solving via instantons [12] the long standing problem of the missing [13, 14, 15, 16, 17, 18, 19, 20, 21] up/down quark mass $\mathbf{10} \mathbf{10} \mathbf{5}^H$ coupling of SU(5)/flipped SU(5) GUTS (from orientifolds of type II compactifications). The latter problem have been also been shown to be solvable by a combination of condensates and Higgs fields by a higher order coupling [22].

In this work we describe a new way of generating the

missing $\mathbf{10} \mathbf{10} \mathbf{5}^H$ in orientifolds of type IIA compactifications by utilizing the formation of strongly coupled condensates extending the work of [22]. In the first part of the this work, we discuss the generation of down-quark masses in flipped SU(5)(FSU(5)). In the second part we generate the up-quark masses in SU(5) GUTS via the same mechanism.

II. THE MISSING YUKAWA COUPLING TERM

A. FLIPPED SU(5) MODELS

Grand unified theories giving rise to SU(5) type gauge group structures [13, 14, 15, 16, 17, 18, 19, 20, 21] arising from 4D compactifications of type II theories on N=1 supersymmetric intersecting brane orientifolds are missing the Yukawa coupling $\mathbf{10} \mathbf{10} \mathbf{5}^H$, which is responsible for generating masses for the down quarks [15, 16, 17, 18, 19]). In an SU(5) GUT the same coupling gives rise to a missing up-quark mass (e.g. see [14, 16, 17, 20, 21]).

In [22], a candidate solution for a perturbative term to the mass of the up-quarks in SU(5) GUTs was suggested where string models with gauge groups in the form $SU(5) \times U(1)^n$ with $n \in \mathbb{Z}$ were used. We have argued that the form of the higher dimensional coupling that gives a mass to the missing $\mathbf{10} \mathbf{10} \mathbf{5}^H$ SU(5) Yukawa's appears from an operator in the form $\frac{1}{M_s^5} \mathbf{10} \mathbf{10} \bar{\mathbf{5}} \bar{\mathbf{5}} \bar{\mathbf{5}} \bar{\mathbf{5}} 1^H 1^H$ where the 1^H gauge singlets are used to cancel the surplus bifundamental charge. In this case, the up-quark mass may come from the condensation of $\bar{\mathbf{5}}$ fermions. Lets us consider the field theory operator $\mathbf{10} \mathbf{10} \bar{\mathbf{5}} \bar{\mathbf{5}} \bar{\mathbf{5}} \bar{\mathbf{5}}$ and rewriting this term in SU(5) notation as

$$\frac{1}{M_s^5} \epsilon^{ijklm} \mathbf{10}^{ij} \mathbf{10}^{kl} \epsilon^{m\phi\chi\psi\omega} \bar{\mathbf{5}}^\phi \bar{\mathbf{5}}^\chi \bar{\mathbf{5}}^\psi \bar{\mathbf{5}}^\omega, \quad (1)$$

we conclude that in components eq. (1) reads

$$\mathbf{10}^{12} \mathbf{10}^{34} \bar{\mathbf{5}}^1 \bar{\mathbf{5}}^2 \bar{\mathbf{5}}^3 \bar{\mathbf{5}}^4 \equiv (u u_L^c)(d_L^c d_L^c d_L^c e_L). \quad (2)$$

*Electronic address: kokorelis@inp.demokritos.gr

†Electronic address: eric@physics.tamu.edu

‡Electronic address: dimitri@physics.tamu.edu

In [22] it was argued that, in the context of field theory (2), can only generate a mass as long as the fermions of the Higgs-like $\bar{5}$ -plets condense. In this work, we will suggest that an alternative explanation for the appearance for such a fermion condensate exists, within the context of SU(5) models constructed on Type IIA orientifolds with intersecting D6-branes where the SU(5) gauge symmetry is extended by Sp(N) gauge groups. The binding mechanism is easy to understand in terms of the gauge forces acting on the fermion condensate as the $\bar{5}$ condense and are charged under Sp(2). The order of the condensate is $\langle (\bar{5}, 1, 1, 2)^4 \rangle = \Lambda^6$, where Λ is the scale the fermion condensate forms. The mass of the up-quarks is determined from (2) and for the present model it reads

$$M_{quarks}^U \approx \frac{\Lambda^6}{M_s^5}. \quad (3)$$

Let us also consider the flipped SU(5) model [20] FSU(5)-IV on Type IIA T6/($Z_2 \times Z_2$) orientifold that also includes metric, NSNS, and RR fluxes [24], [25]. In this flipped SU(5) model the complete gauge symmetry is $U(5) \times U(1)^{10} \times USp(10)^2 \times USp(8) \times USp(2)$.

stk	N	$(n_1, l_1)(n_2, l_2)(n_3, l_3)$	A	S	O64
a	10	(1, 3)(1, 1)(0, -1)	2	-2	-1
b	2	(1, -3)(0, -1)(-1, 1)	-2	2	0
c	2	(1, 3)(-1, 1)(-1, 0)	2	-2	0
d	2	(1, 1)(-1, -3)(0, 1)	2	-2	-1
e	2	(-5, 9)(-5, -3)(1, 0)	-30	30	0
f	2	(2, 0)(-1, 3)(-1, -3)	0	0	6
g	2	(1, -9)(-1, 0)(-1, -3)	-6	6	3
h	2	(1, -7)(0, 1)(7, -3)	0	0	0
i	2	(0, 2)(4, -3)(3, -4)	0	0	0
j	2	(1, -3)(-1, 0)(-1, -1)	-2	2	1
k	2	(0, 2)(-3, -1)(1, 3)	0	0	0
O6 ¹	10	(2, 0)(1, 0)(1, 0)	-	-	-
O6 ²	10	(2, 0)(0, -1)(0, 1)	-	-	-
O6 ³	8	(0, -2)(1, 0)(0, 1)	-	-	-
O6 ⁴	2	(0, -2)(0, 1)(1, 0)	-	-	-

TABLE I: D6-brane configurations and intersection numbers (Part 1) for the Model FSU(5)-IV on Type IIA $\mathbf{T}^6/(\mathbb{Z}_2 \times \mathbb{Z}_2)$ orientifold with fluxes. The complex structure parameters are $\chi_1 = 2/3$, $\chi_2 = 1$, and $\chi_3 = 1$. We have listed only the intersections of the branes with the Sp(2) generating brane.

We observe that only the Yukawa mass coupling term giving a mass to the up quarks is allowed (there is no obvious mass term for the leptons as well)

$$\langle \mathbf{10}_{(2,0^{10})} \bar{\mathbf{5}}_{(-1,1,0^9)} \bar{\mathbf{5}}_{(-1,-1,0^9)}^H \rangle, \langle \bar{\mathbf{5}}_{(-1,1,0^9)} \mathbf{1}_{(0,2,0^9)} \mathbf{5}_{(1,1,0^9)}^H \rangle \quad (4)$$

We used for convenience the notation $k^n \equiv (k, k, \dots, k)$ with k repeated n times. Also as there is no mass term for the down-quarks of the form $\mathbf{10}_{(2,0^{10})} \cdot \mathbf{10}_{(2,0^{10})} \cdot \mathbf{5}_{(1,1,0^9)}^H$ as this is not allowed by $U(1)_a, U(1)_b$ charge conservation. This is the usual problem of any SU(5)/flipped

SU(5) GUT from any type II orientifold compactifications e.g.[14, 15, 16, 17, 18, 19]. The down-quark masses may be generated by the expression

$$\frac{1}{M_s^5} ((10, 1^{10}; 1^3, 1)_{(2,0^{10})})^2 \cdot \left((\bar{\mathbf{5}}, 1^{10}; 1^3, 2)_{(-1,0^{10})}^H \right)^4. \quad (5)$$

In order to show that the $\bar{5}$'s condense we will adjoint split the gauge group $U(5) \rightarrow U(3) \times U(2)$, so that the U(5)-D6-brane splits into a and a1 branes, each one associated with the U(3), U(2) brane stacks. The anomaly free hypercharge corresponding to the decomposition of table (II) is

$$Y = \frac{1}{6}U(1)_a - \frac{1}{2}U(1)_b + \frac{1}{2}U(1)_c + \frac{1}{2}U(1)_d - \frac{1}{2}U(1)_j + \frac{1}{2}U(1)_e + \frac{1}{2}U(1)_f - \frac{1}{2}U(1)_g - \frac{1}{2}U(1)_h + \frac{1}{2}U(1)_i - \frac{1}{2}U(1)_k \quad (6)$$

We have parametrized the number of non-chiral states

		$SU(3) \times SU(2)_{a_1} \times U(1)_b \cdots \times USp(2)$	
Intersection	Multiplicity		Y
a,2	1	$(\bar{3}, 1, 1^{10}; 1, 1, 1, 2)_{(-1,0,0^{10})}$	$-\frac{1}{6}$
a ₁ ,2	1	$(1, 2, 1^{10}; 1, 1, 1, 2)_{(0,-1,0^{10})}$	0
b,2	B'	$(1^{12}; 1, 1, 1, 2)_{(0,0,1,0^9)}$	$-\frac{1}{2}$
b,2	B'	$(1^{12}; 1, 1, 1, 2)_{(0,0,-1,0^9)}$	$\frac{1}{2}$
c,2	C'	$(1^{12}; 1, 1, 1, 2)_{(0,0,0,1,0^8)}$	$\frac{1}{2}$
c,2	C'	$(1^{12}; 1, 1, 1, 2)_{(0,0,0,-1,0^8)}$	$-\frac{1}{2}$
d,2	1	$(1^{12}; 1, 1, 1, 2)_{(0^4,-1,0^7)}$	$\frac{1}{6}$
e,2	E'	$(1^{12}; 1, 1, 1, 2)_{(0^5,1,0^6)}$	$\frac{1}{2}$
e,2	E'	$(1^{12}; 1, 1, 1, 2)_{(0^5,-1,0^6)}$	$-\frac{1}{2}$
f,2	6	$(1^{12}; 1, 1, 1, 2)_{(0^6,1,0^5)}$	$\frac{1}{2}$
g,2	3	$(1^{12}; 1, 1, 1, 2)_{(0^7,1,0^4)}$	$-\frac{1}{2}$
h,2	H'	$(1^{12}; 1, 1, 1, 2)_{(0^8,1,0^3)}$	$-\frac{1}{2}$
h,2	H'	$(1^{12}; 1, 1, 1, 2)_{(0^8,-1,0^3)}$	$\frac{1}{2}$
i,2	I'	$(1^{12}; 1, 1, 1, 2)_{(0^9,1,0^2)}$	$\frac{1}{2}$
i,2	I'	$(1^{12}; 1, 1, 1, 2)_{(0^9,-1,0^2)}$	$-\frac{1}{2}$
j,2	1	$(1^{12}; 1, 1, 1, 2)_{(0^{10},1,0)}$	$-\frac{1}{2}$
k,2	K'	$(1^{12}; 1, 1, 1, 2)_{(0^{11},1)}$	$-\frac{1}{2}$
k,2	K'	$(1^{12}; 1, 1, 1, 2)_{(0^{11},-1)}$	$\frac{1}{2}$

TABLE II: N=1 multiplets charged under Sp(2) in FSU(5). The gauge group is $SU(3) \times SU(2)_{a_1} \times U(1)_b \times U(1)_c \times U(1)_d \times U(1)_e \times U(1)_f \times U(1)_g \times U(1)_h \times U(1)_i \times U(1)_j \times U(1)_k \times USp(10)^2 \times USp(8) \times USp(2)$.

in intersections where branes are parallel in at last one tori by B', C', E', H', I', K' . The β -function for Sp(2) is $b^{Sp(2)} = 8 + B' + C' + E' + H' + I' + K' - 6$ and may become negative as long the number of non-chiral states is either 0 or if they decouple at a typical mass of order M_s , higher than the condensation scale Λ_{cond} . We may also need to give a mass Λ_{cond} to the multiplets from the

intersections (a,2), (a₁,2), (f,2), (g,2). The leptons may get a mass from

$$\frac{1}{M_s^6} \bar{\mathbf{5}}_{(-1,1,0^9)} \mathbf{1}_{(0,2,0^9)} \mathbf{5}_{(1,1,0^9)}^H \langle (1^{12}; 1, 1, 1, 2)_{(0,-1,0^{10})} \rangle^4 \quad (7)$$

where the states $\langle \dots \rangle$ from the intersection (a₁,2) may condense and decouple. In Table III, using eq.(3), we exhibit the condensation scale against the string scale given the phenomenological down-quark masses; $3.5 < m_d < 6$ MeV, $M_s \approx 105$ MeV, $M_b \approx 4.20$ GeV [23]. Thus

GeV		$M_S = 10^{16}$	$M_S = 10^{17}$	$M_S = 10^{18}$
Quark	M_{quark}			
d	$5 \cdot 10^{-3}$	$8.9 \cdot 10^{12}$	$3.25 \cdot 10^{14}$	$4.1 \cdot 10^{14}$
s	$105 \cdot 10^{-3}$	$1.5 \cdot 10^{13}$	$1.0 \cdot 10^{14}$	$6.9 \cdot 10^{14}$
b	4.2	$2.7 \cdot 10^{13}$	$1.9 \cdot 10^{14}$	$1.3 \cdot 10^{15}$

TABLE III: Λ_{cond} vs down-quark masses using eq.(3).

a generic FSU(5) string model may therefore describe the masses of the down-quarks as long as its condensation scale is in the range

$$10^{12} \text{ GeV} < \Lambda_{FSU(5)}^{cond} < 10^{15} \text{ GeV} \quad (8)$$

If we consider the running gauge Sp(2) coupling following [7] with $b_{Sp(2)} = -2$ where we have set, B', C', E', H', I', K' to zero and the states from intersections (a,2), (a₁,2), (g,2) decouple, Λ_{cond} is determined in table (IX). M_s can be higher than 10^{18} GeV since its value depends on the number of non-chiral states which we will not determine here. Values of Λ_{cond} higher from Λ_{run}^{cond} may be

	$M_S = 10^{16}$	$M_S = 10^{18}$	$M_S = 10^{19}$	$M_S = 10^{21}$
Λ_{run}^{cond}	$2.6 \cdot 10^{10}$	$2.6 \cdot 10^{12}$	$2.6 \cdot 10^{13}$	$2.6 \cdot 10^{14}$

TABLE IV: Λ_{run}^{cond} from the running Sp(2) gauge coupling.

understood (from the exact string amplitude of the associated fermionic correlator of eq.(2)) which involves a suppression factor lowering Λ_{cond} to Λ_{run}^{cond} .

III. SU(5) GUTS

Let us consider for example the SU(5)-type D6-brane model II.1.4. of [27]. It requires three stacks a , c and d of D6-branes giving rise to a $U(5)_a \times U(1)_c \times U(1)_d$ gauge symmetry intersecting at angles in IIA orientifolds of $Z_2 \times Z_2$ toroidal compactifications [26]. The $U(5)_a$ splits into $SU(5)_a \times U(1)_a$, where the anomalous $U(1)_a$ gauge boson becomes massive via the generalized Green-Schwarz mechanism and $U(1)_a$ appears as a global symmetry in the effective action. The matter transforming as $\mathbf{10}$ under $SU(5)_a$ arises at intersections of stack a with its image a' ; the matter fields transforming as $\bar{\mathbf{5}}$ as well

as Higgs fields $\mathbf{5}_H$ and $\bar{\mathbf{5}}_H$ are located at intersections of stack a with c and c' or d and d' . The key input in the construction of the D-brane model is summarized in Tables V and VI. Table VI lists the candidate Higgs fields.

	$U(5) \times U(1)_c \times U(1)_d \times USp(2)$								
stack	N	$(n^1, l^1)(n^2, l^2)(n^3, l^3)$	$n_{\square\square}$	$n_{\square}$	c	d	c'	d'	2
a	10	$(0, -1)(1, 4)(1, 1)$	-3	3	0	18	0	8	-1
c	2	$(-1, 3)(-1, 4)(1, 1)$	-6	-42	-	-18	-	-72	-3
d	2	$(-1, 0)(-1, 2)(7, 1)$	13	-13	-	-	-	-	0
2	2	$(1, 0) \times (0, 1) \times (0, -2)$	$7x_B = 28x_A = 20x_C$						

TABLE V: Wrapping numbers of D6-branes.

sector	number	$U(5)_a \times U(1)_c \times U(1)_d \times USp(2)$
(a, a')	3	$(10, 1, 1, 1)_{(2,0,0)} + (\bar{15}, 1, 1, 1)_{(-2,0,0)}$
(a, c)	0	$(5, 1, 1, 1)_{(1,-1,0)} + (\bar{5}, 1, 1, 1)_{(-1,1,0)}$
(a, c')	0	$(\bar{5}, 1, 1, 1)_{(-1,-1,0)}^H + (5, 1, 1, 1)_{(1,1,0)}^H$
$(a, 2)$	-1	$(\bar{5}, 1, 1, 2)_{(-1,0,0)}^H$

TABLE VI: N=1 multiplet matter spectrum for $SU(5)$.

As can be seen from tables (V) and (VI), the only mass terms allowed are the ones that are associated with the Yukawa couplings giving masses to the down quarks and leptons respectively

$$\langle \mathbf{10}_{(2,0,0)} \bar{\mathbf{5}}_{(-1,1,0)} \bar{\mathbf{5}}_{(-1,-1,0)}^H \rangle, \langle \bar{\mathbf{5}}_{(-1,1,0)} \mathbf{1}_{(0,-2,0)} \mathbf{5}_{(1,1,0)}^H \rangle \quad (9)$$

and there is no mass term for the up-quarks of the form

$$\mathbf{10}_{(2,0,0)} \cdot \mathbf{10}_{(2,0,0)} \cdot \mathbf{5}_{(1,1,0)}^H \quad (10)$$

as this is not allowed by $U(1)_a, U(1)_b$ charge conservation. This is the usual problem of any SU(5) GUT from any type II orientifold compactifications e.g.[14, 15, 16, 17]. Thus, working without loss of generality in the context of SU(5) model shown in Tables (V) and (VI), a perturbative mass term for the up-quark masses of the form

$$\frac{1}{M_s^5} ((10, 1, 1, 1)_{(2,0,0)})^2 \cdot ((\bar{5}, 1, 1, 2)_{(-1,0,0)}^H)^4 \quad (11)$$

exists. In order to demonstrate that the Sp(2) gauge group will become strongly coupled, we will use adjoint breaking (AB) to break the SU(5) down to the $SU(3)_a \times SU(2)_b \times U(1)_Y$, $Y = (1/6)U(1)_a + (1/2)U(1)_b - (1/2)U(1)_c + (1/2)U(1)_d$, which is equivalent to splitting the stacks on one torus. To establish notation, $U(1)_a, U(1)_b$ are the U(1)'s within U(3), U(2) (of U(5)). Thus, we only present explicitly the states charged under Sp(2) in table (VII). We have parametrized the appearance of an arbitrary number of non-chiral states becoming massless by A . Since $b_a^{Sp(2)} = A - 2$, the b -function becomes asymptotically free when the number of non-chiral multiplets becomes either 0 or 1.

$SU(3) \times SU(2)_w \times U(1)_c \times U(1)_d \times USp(2)$		
Multiplicity		Y
1	$(\bar{3}, 1, 1, 1, 2)_{(-1,0,0,0)}$	$-\frac{1}{6}$
1	$(1, 2, 1, 1, 2)_{(0,-1,0,0)}$	$-\frac{1}{2}$
3	$(1, 1, 1, 1, 2)_{(0,0,-1,0)}$	$\frac{1}{2}$
A	$(1, 1, 1, 1, \bar{2})_{(0,0,0,1)}$	$\frac{1}{2}$
A	$(1, 1, 1, 1, 2)_{(0,0,0,-1)}$	$-\frac{1}{2}$

TABLE VII: N=1 multiplets charged under Sp(2).

In Table VIII, utilizing the relation (3), we show how the condensation scale depends on the string scale given the values of the phenomenologically determined up-quark masses with $m_u = 1.5$ to 3.3 MeV; $m_c = 1.27$ GeV, $m_t \approx 171$ GeV [23].

Quark	GeV	$M_S = 10^{16}$	$M_S = 10^{17}$	$M_S = 10^{18}$
c	1.27	$2.2 \cdot 10^{13}$	$1.5 \cdot 10^{14}$	$1.0 \cdot 10^{15}$
u	$2.5 \cdot 10^{-3}$	$7.9 \cdot 10^{12}$	$5.4 \cdot 10^{13}$	$3.7 \cdot 10^{14}$
t	171.3	$5.1 \cdot 10^{13}$	$3.4 \cdot 10^{14}$	$2.3 \cdot 10^{15}$

TABLE VIII: Λ_{cond} (GeV) vs up-quark masses *using eq.(3)*.

Thus a generic SU(5) string model may therefore describe the masses of the up-quarks as long as its condensation scale is in the range

$$10^{12} \text{ GeV} < \Lambda_{SU(5)}^{cond} < 10^{15} \text{ GeV} \quad (12)$$

The range of Λ_{run}^{cond} is what one gets from the strong coupling of Sp is similar to that of FSU(5)(we have set $A = 0$, $\chi_C = 28/20$).

	$M_S = 10^{16}$	$M_S = 10^{18}$	$M_S = 10^{19}$	$M_S = 10^{21}$
Λ_{run}^{cond}	$2.4 \cdot 10^9$	$2.4 \cdot 10^{11}$	$2.4 \cdot 10^{12}$	$2.4 \cdot 10^{14}$

TABLE IX: Λ_{run}^{cond} from the running Sp(2) gauge coupling.

We have presented a new perturbative mechanism to generate the masses of the up/down quarks through fermion condensates within N=1 supersymmetric SU(5)/flipped SU(5) constructions from intersecting D6-branes. We have also demonstrated that as Λ^{cond} appears to be in the same range as that expected from the running of Sp Λ_{run}^{cond} for up and down quarks (eqn's (8),(12)), this indicates possible flavour independence. Also, from the Λ^6 dependence on the condensation scale, expected small differences in condensate values naturally may provide us with the fermion mass textures. It would be also important to further study this issue by calculating the relevant string amplitudes. We have also seen that a necessary condition for the existence of the fermion condensate giving a mass to the missing quark masses, is to simultaneously demand that the number of non-chiral states charged under the Sp(N) condensing gauge group may be fixed so that Sp(N) may have negative β -function.

Acknowledgements: C.K wishes to thank E.G.Floratos for discussions. The work of D.V.N. is supported by DOE grant DEFG03-95-Er-40917.

-
- [1] G. 't Hooft, Phys. Rev. D **14** (1976) 3432 [Erratum-ibid. D **18** (1978) 2199].
- [2] E. Witten, Nucl. Phys. B **149** (1979) 285.
- [3] M. Dine, N. Seiberg, X. G. Wen and E. Witten, Nucl. Phys. B **278** (1986) 769.
- [4] D. Lust and T. R. Taylor, Phys. Lett. B **253** (1991) 335.
- [5] R. Blumenhagen, M. Cvetič, P. Langacker and G. Shiu, Ann. Rev. Nucl. Part. Sci. **55** (2005) 71
- [6] R. Blumenhagen, B. K rs, D. L st and S. Stieberger, Phys. Rept. **445**, 1 (2007) [arXiv:hep-th/0610327].
- [7] M. Cvetič, P. Langacker and G. Shiu, Phys. Rev. D **66** (2002) 066004 [arXiv:hep-ph/0205252].
- [8] I. Antoniadis, J. R. Ellis, J. S. Hagelin and D. V. Nanopoulos, Phys. Lett. B **231** (1989) 65; Phys. Lett. B **194** (1987) 231.
- [9] R. Blumenhagen, M. Cvetič, S. Kachru and T. Weigand, arXiv:0902.3251 [hep-th].
- [10] L. E. Ibanez and R. 2. Richter, JHEP **0903** (2009) 090
- [11] M. Cvetič, J. Halverson and R. 2. Richter, arXiv:0905.3379 [hep-th].
- [12] R. Blumenhagen, M. Cvetič, D. L st, R. Richter and T. Weigand, Phys. Rev. Lett. **100**, 061602 (2008)
- [13] C. Bachas, arXiv:hep-th/9503030.
- [14] R. Blumenhagen, B. K rs, D. Lust and T. Ott, Nucl. Phys. B **616** (2001) 3 [arXiv:hep-th/0107138].
- [15] J.R. Ellis, P. Kanti and D. V. Nanopoulos, Nucl. Phys. B **647** (2002) 235 [arXiv:hep-th/0206087].
- [16] M. Axenides, E. Floratos and C. Kokorelis, JHEP **0310** (2003) 006 [arXiv:hep-th/0307255].
- [17] C. Kokorelis, Nucl. Phys. B **732** (2006) 341
- [18] C. M. Chen, G. V. Kraniotis, V. E. Mayes, D. V. Nanopoulos and J. W. Walker, Phys. Lett. B **611** (2005) 156 Phys. Lett. B **625** (2005) 96
- [19] C.M. Chen, V. E. Mayes and D. V. Nanopoulos, Phys. Lett. B **633** (2006) 618 [arXiv:hep-th/0511135].
- [20] C.M. Chen, T. Li and D. V. Nanopoulos, Nucl. Phys. B **751** (2006) 260 [arXiv:hep-th/0604107].
- [21] C. Bachas, M. Bianchi, R. Blumenhagen, D. Lust and T. Weigand, JHEP **0808**, 016 (2008)
- [22] C. Kokorelis, arXiv:0812.4804 [hep-th].
- [23] C. Amsler et al., Physics Letters B667, 1 (2008)
- [24] G. Villadoro and F. Zwirner, JHEP **0506** (2005) 047
- [25] P. G. Camara, A. Font and L. E. Ibanez, JHEP **0509**, 013 (2005) [arXiv:hep-th/0506066].
- [26] M. Cvetič, G. Shiu and A. M. Uranga, Nucl. Phys. B **615**, 3 (2001) [arXiv:hep-th/0107166].
- [27] M. Cvetič, I. Papadimitriou and G. Shiu, Nucl. Phys. B **659** (2003) 193 [Erratum-ibid. B **696** (2004) 298]
- [28] M. Cvetič and I. Papadimitriou, Phys. Rev. D **68**, 046001 (2003) [Erratum-ibid. D **70**, 029903 (2004)]